



# $\nu_e$ and $\bar{\nu}_e$ disappearance in Gallium and reactor experiments [1]

MARIO A. ACERO <sup>(a,b,c)</sup>, CARLO GIUNTI <sup>(b)</sup>, MARCO LAVEDER <sup>(d)</sup>

(a) Dipartimento di Fisica Teorica, Università di Torino. (b) INFN, Sezione di Torino.  
(c) Laboratoire d'Annecy-le-Vieux de Physique Théorique LAPTH, Université de Savoie, CNRS/IN2P3.  
(d) Dipartimento di Fisica "G. Galilei", Università di Padova, and INFN, Sezione di Padova.



## ABSTRACT

The disappearance of electron neutrinos observed in the Gallium radioactive source experiments is analyzed in the effective framework of two-neutrino mixing. We found an indication of neutrino disappearance due to neutrino oscillations with  $\sin^2 2\vartheta \gtrsim 0.03$  and  $\Delta m^2 \gtrsim 0.1 \text{ eV}^2$ . We study the compatibility of this result with the data of the Bugey and Chooz reactor short-baseline antineutrino disappearance experiments, founding an indication in favor of neutrino oscillations with  $0.01 \lesssim \sin^2 2\vartheta \lesssim 0.07$  and  $1.8 \text{ eV}^2 \lesssim \Delta m^2 \lesssim 1.9 \text{ eV}^2$ , from the Bugey data, which is compatible with the Gallium allowed region of the mixing parameters. This indication persists in the combined analyses of Gallium, Bugey, and Chooz data.

## INTRODUCTION

Solar, atmospheric, reactor and accelerator neutrino experiments give very robust evidence of three-neutrino mixing [2, 3]. However, data from LSND, MiniBooNE (at low energy) and the Gallium radioactive source experiments show some anomalies which open a window to the possible existence of exotic neutrino physics beyond the three-neutrino mixing. The Gallium radioactive source experiments (GALLEX [4] and SAGE [5, 6]) measured a lower electron neutrino flux than the expected one, and this can be interpreted as an indication of the disappearance of electron neutrinos due to neutrino oscillations.

## Gallium experiments

The Gallium radioactive source experiment consist in the detection of electron neutrinos produced by artificial  $^{51}\text{Cr}$  and  $^{37}\text{Ar}$  radioactive sources which decay through electron capture



The neutrinos are detected through the reaction  $\nu_e + ^{71}\text{Ga} \rightarrow ^{71}\text{Ge} + e^-$ . We present the results of the fit of the data of Gallium radioactive source experiments in terms of effective two-neutrino oscillations. The survival probability of electron (anti)neutrinos with energy  $E$  at a distance  $L$  from the source is given by


$$P_{\nu_e \rightarrow \nu_e}^{(-)}(L, E) = 1 - \sin^2 2\vartheta \sin^2 \left( \frac{\Delta m^2 L}{4E} \right), \quad (1)$$

where  $\vartheta$  is the mixing angle and  $\Delta m^2$  is the squared-mass difference (see Refs. [2, 3]). We use the theoretical value of the ratio  $R$  of the predicted  $^{71}\text{Ge}$  production rates in each of the Gallium radioactive source experiments in the cases of presence and absence of neutrino oscillations given by

$$R = \frac{\int dV L^{-2} \sum_i (\text{B.R.})_i \sigma_i P_{\nu_e \rightarrow \nu_e}(L, E_i)}{\sum_i (\text{B.R.})_i \sigma_i \int dV L^{-2}}, \quad (2)$$

where  $i$  is the index of the  $\nu_e$  lines emitted in  $^{51}\text{Cr}$  or  $^{37}\text{Ar}$ .

The result of the combined least-squares analysis of the four Gallium source experiments is shown in Fig. (1). One can see that there is an allowed region in the  $\sin^2 2\vartheta - \Delta m^2$  plane at  $1\sigma$  for  $\Delta m^2 \gtrsim 0.6 \text{ eV}^2$  and  $0.08 \lesssim \sin^2 2\vartheta \lesssim 0.4$ .



**Figure 1.** (Left) Allowed region in the oscillation parameter space for the individual Gallium radioactive source experiments. (Right) Allowed regions in the oscillation parameter space and marginal  $\Delta\chi^2$ 's parameters for the combined fit of the results of the four radioactive source experiments.

	Ga	Bu	Ga+Bu	Bu+Ch	Ga+Ch	Ga+Bu+Ch
$\chi^2_{\min}$	2.69	46.55	52.59	47.12	6.57	53.40
NDF	2	53	57	54	3	58
GoF	0.26	0.72	0.64	0.73	0.087	0.65
$\sin^2 2\vartheta_{\text{bf}}$	0.23	0.043	0.057	0.036	0.079	0.05
$\Delta m^2_{\text{bf}} [\text{eV}^2]$	2.00	1.85	1.85	1.85	1.73	1.85

**Table 1.** Values of  $\chi^2_{\min}$ , number of degrees of freedom (NDF) and best-fit values of  $\sin^2 2\vartheta$  and  $\Delta m^2$  from the fit of different combinations of the results of the Gallium radioactive source experiments and the Bugey and Chooz reactor experiments.

## Bugey and Chooz reactor experiments


Reactor neutrino experiments detect antineutrinos through the reaction  $\bar{\nu}_e + p \rightarrow n + e^+$ . In this process, the neutrino energy is related with the positron energy by  $E_\nu = E_{e^+} + 1.8 \text{ MeV}$ . The Bugey experiment used three source-detector distances ( $L = 15, 40, 95 \text{ m}$ ), while in Chooz, used a distance about 1 km. For the Bugey experiment we use the ratio of observed and expected (in the case of no oscillation) positron spectra given in Fig. 17 of Ref. [7], in which there are  $N_j = 25, 25, 10$  energy bins. We analyze the data with the following  $\chi^2$ :

$$\chi^2 = \sum_{j=1}^3 \left\{ \sum_{i=1}^{N_j} \frac{[(A a_j + b(E_{ji} - E_0)) R_{ji}^{\text{the}} - R_{ji}^{\text{exp}}]^2}{\sigma_{a_j}^2} + \frac{(a_j - 1)^2}{\sigma_{a_j}^2} \right\} + \frac{(A - 1)^2}{\sigma_A^2} + \frac{b^2}{\sigma_b^2}, \quad (3)$$

where  $E_{ji}$  is the central energy of the  $i$ th bin in the positron kinetic energy spectrum measured at the  $L_j$  source-detector distance,  $R_{ji}^{\text{exp}}$  is the measured ratio and

$$R_{ji}^{\text{the}} = \frac{\int dL L^{-2} \int_{E_{ji}-\Delta E_j/2}^{E_{ji}+\Delta E_j/2} dE \int_{-\infty}^{+\infty} dT_e F(E, T_e) P_{\bar{\nu}_e \rightarrow \bar{\nu}_e}(L, E_\nu)}{\Delta E_j \int dL L^{-2}}. \quad (4)$$

In the case of the Chooz experiment, this gives constraints on  $\sin^2 2\vartheta$  for  $\Delta m^2 \gtrsim 10^{-3} \text{ eV}^2$ , so for our purpose, the Chooz experiment is only sensitive to the average survival probability  $\langle P_{\nu_e \rightarrow \nu_e}^{(-)} \rangle = 1 - \frac{1}{2} \sin^2 2\vartheta$ .



**Figure 2.** (Up) Allowed region in the oscillation parameter space and histograms with the Best Fit obtained from for the Bugey reactor experiment (Up-Right). (Down-Left) Allowed regions in the oscillation parameter space for the combined fit of the Bugey and Chooz reactor experiments. (Down-Right) Allowed regions for the combined fit of the Gallium radioactive source experiments and the Bugey and Chooz reactor experiments.

## Conclusions

In the framework of two-neutrino mixing, we found that, from the analysis of the Gallium radioactive source experiments, there is an indication of electron neutrino disappearance due to neutrino oscillations with  $\sin^2 2\vartheta \gtrsim 0.03$  and  $\Delta m^2 \gtrsim 0.1 \text{ eV}^2$ . This result is compatible with the data from Bugey and Chooz reactor experiments. Besides, the Bugey data present an indication in favor of neutrino oscillations with  $0.01 \lesssim \sin^2 2\vartheta \lesssim 0.07$  and  $1.8 \text{ eV}^2 \Delta m^2 \simeq 1.9 \text{ eV}^2$ . Such a disappearance of electron neutrinos due to  $\Delta m^2 \gtrsim 0.1 \text{ eV}^2$  is an indication of the possible existence of at least one light sterile neutrino with a mass of the order about 1 eV [3].

## References

- [1] Mario A. Acero, Carlo Giunti and Marco Laveder, arXiv:0711.4222.
- [2] S.M. Bilenky and S.T. Petcov, Rev. Mod. Phys. 59 (1987) 671.
- [3] C. Giunti and C.W. Kim, Fundamentals of Neutrino Physics and Astrophysics (Oxford University Press, 2007).
- [4] GALLEX, W. Hampel et al., Phys. Lett. B420 (1998) 114.
- [5] SAGE, J.N. Abdurashitov et al., Phys. Rev. C59 (1999) 2246, arXiv:hep-ph/9803418.
- [6] J.N. Abdurashitov et al., Astropart. Phys. 25 (2006) 349, arXiv:nucl-ex/0509031.
- [7] Bugey, B. Achkar et al., Nucl. Phys. B434 (1995) 503.