

Review of Present Hints and Possible Combined Analysis

Carlo Giunti

INFN, Sezione di Torino, and Dipartimento di Fisica Teorica, Università di Torino

<mailto://giunti@to.infn.it>

Neutrino Unbound: <http://www.nu.to.infn.it>


BEYOND3NU

Workshop on Beyond Three Family Neutrino Oscillations

3-4 May 2011, LNGS

Collaboration with Marco Laveder (Padova University)

Three-Neutrino Mixing Paradigm



Two scales of $\Delta m^2 \iff$ Three-Neutrino Mixing

$$\Delta m_{\text{SOL}}^2 = \Delta m_{21}^2 \simeq 7.6 \times 10^{-5} \text{ eV}^2$$

$$\Delta m_{\text{ATM}}^2 \simeq |\Delta m_{31}^2| \simeq |\Delta m_{32}^2| \simeq 2.4 \times 10^{-3} \text{ eV}^2$$



LSND

[LSND, PRL 75 (1995) 2650; PRC 54 (1996) 2685; PRL 77 (1996) 3082; PRD 64 (2001) 112007]

$$\bar{\nu}_\mu \rightarrow \bar{\nu}_e$$

$$L \simeq 30 \text{ m}$$

$$20 \text{ MeV} \leq E \leq 200 \text{ MeV}$$



$$\Delta m_{\text{LSND}}^2 \gtrsim 0.2 \text{ eV}^2 \quad (\gg \Delta m_{\text{ATM}}^2 \gg \Delta m_{\text{SOL}}^2)$$



MiniBooNE Antineutrinos

[PRL 103 (2009) 111801; PRL 105 (2010) 181801]

$$\bar{\nu}_\mu \rightarrow \bar{\nu}_e$$

$L \simeq 541 \text{ m}$

$475 \text{ MeV} \leq E \lesssim 3 \text{ GeV}$




[MiniBooNE, PRL 105 (2010) 181801, arXiv:1007.1150]

Agreement with LSND $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ signal!

[talk by M. Sorel]

Similar L/E but different L and $E \rightarrow$ Oscillations!

Beyond Three-Neutrino Mixing



Standard Model

- ▶ Neutrinos are the only massless fermions
- ▶ Neutrinos are the only fermions with only left-handed component ν_L

Extension of the SM: Massive Neutrinos

- ▶ Simplest extension: introduce right-handed component ν_R
- ▶ Dirac mass $m_D \bar{\nu}_R \nu_L +$ Majorana mass $m_M \bar{\nu}_R^c \nu_R$
- ▶ $\nu_{eL}, \nu_{\mu L}, \nu_{\tau L} + \nu_{eR}, \nu_{\mu R}, \nu_{\tau R} \implies 6$ massive Majorana neutrinos

Sterile Neutrinos

- ▶ Light anti- ν_R are called sterile neutrinos

$$\nu_R^c \rightarrow \nu_s \quad (\text{left-handed})$$

- ▶ Sterile means no standard model interactions
- ▶ Active neutrinos (ν_e, ν_μ, ν_τ) can oscillate into sterile neutrinos (ν_s)
- ▶ Observables:
 - ▶ Disappearance of active neutrinos
 - ▶ Indirect evidence through combined fit of data (current indication)
- ▶ Powerful window on new physics beyond the Standard Model

How many Sterile Neutrinos?


$$e^+ e^- \rightarrow Z \xrightarrow{\text{invisible}} \sum_{a=\text{active}} \nu_a \bar{\nu}_a$$

$$N_{\nu_{\text{active}}} = 2.9840 \pm 0.0082$$

[LEP, Phys. Rept. 427 (2006) 257, hep-ex/0509008]

3 light active flavor neutrinos

$$\nu_e \quad \nu_\mu \quad \nu_\tau$$



mixing $\Rightarrow \nu_{\alpha L} = \sum_{k=1}^N U_{\alpha k} \nu_{kL} \quad \alpha = e, \mu, \tau$ $N \geq 3$
no upper limit!


Mass Basis: $\nu_1 \quad \nu_2 \quad \nu_3 \quad \nu_4 \quad \nu_5 \quad \dots$

Flavor Basis: $\nu_e \quad \nu_\mu \quad \nu_\tau \quad \nu_{s_1} \quad \nu_{s_2} \quad \dots$

ACTIVE STERILE

Cosmology

- N_s = number of thermalized sterile neutrinos (not necessarily integer)



- CMB and LSS in Λ CDM:

[Hamann, Hannestad, Raffelt, Tamborra, Wong, PRL 105 (2010) 181301, arXiv:1006.5276]

$$N_s = 1.61 \pm 0.92 \quad m_{\nu_s} < 0.70 \text{ eV} \quad (95\% \text{ C.L.}) \quad [\text{talk by A. Melchiorri}]$$

[Giusarma, Corsi, Archidiacono, de Putter, Melchiorri, Mena, Pandolfi, arXiv:1102.4774]

$$N_s = 0.22 \pm 0.59 \quad [\text{Cyburt, Fields, Olive, Skillman, AP 23 (2005) 313, astro-ph/0408033}]$$

- BBN:

$$N_s = 0.64^{+0.40}_{-0.35} \quad [\text{Izotov, Thuan, ApJL 710 (2010) L67, arXiv:1001.4440}]$$

SBL Oscillation Probabilities in 3+1 Schemes

$$P_{\nu_\alpha \rightarrow \nu_\beta} = \sin^2 2\vartheta_{\alpha\beta} \sin^2 \left(\frac{\Delta m^2 L}{4E} \right)$$

$$\sin^2 2\vartheta_{\alpha\beta} = 4|U_{\alpha 4}|^2 |U_{\beta 4}|^2$$

No CP Violation!

$$P_{\nu_\alpha \rightarrow \nu_\alpha} = 1 - \sin^2 2\vartheta_{\alpha\alpha} \sin^2 \left(\frac{\Delta m^2 L}{4E} \right)$$

$$\sin^2 2\vartheta_{\alpha\alpha} = 4|U_{\alpha 4}|^2 (1 - |U_{\alpha 4}|^2)$$

Perturbation of 3ν Mixing

$$|U_{e4}|^2 \ll 1, \quad |U_{\mu 4}|^2 \ll 1, \quad |U_{\tau 4}|^2 \ll 1, \quad |U_{s4}|^2 \simeq 1$$


$$U = \begin{pmatrix} U_{e1} & U_{e2} & U_{e3} & U_{e4} \\ U_{\mu 1} & U_{\mu 2} & U_{\mu 3} & U_{\mu 4} \\ U_{\tau 1} & U_{\tau 2} & U_{\tau 3} & U_{\tau 4} \\ U_{s1} & U_{s2} & U_{s3} & U_{s4} \end{pmatrix}$$

SBL

$\sin^2 2\vartheta_{\alpha\alpha} \ll 1$

\Downarrow

$|U_{\alpha 4}|^2 \simeq \frac{\sin^2 2\vartheta_{\alpha\alpha}}{4}$




3+1 Schemes

GoF = 32%


PGoF = 0.89%

- ▶ Tension between LSND + KARMEN + MiniBooNE $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ and MiniBooNE $\nu_\mu \rightarrow \nu_e \implies$ CP Violation?
- ▶ 3+2 \implies CP Violation OK [Sorel, Conrad, Shaevitz, PRD 70 (2004) 073004, hep-ph/0305255; Maltoni, Schwetz, PRD 76, 093005 (2007), arXiv:0705.0107; Karagiorgi et al, PRD 80 (2009) 073001, arXiv:0906.1997]
- ▶ 3+1+NSI \implies CP Violation OK [Akhmedov, Schwetz, JHEP 10 (2010) 115, arXiv:1007.4171]

$\bar{\nu}_e$ Disappearance



ν_μ and $\bar{\nu}_\mu$ Disappearance



New Reactor $\bar{\nu}_e$ Fluxes

[Mueller et al., arXiv:1101.2663]

[Mention et al., arXiv:1101.2755]

[talk by D. Lhuillier]

ATM constraint on $|U_{\mu 4}|^2$

[Maltoni, Schwetz, PRD 76 (2007) 093005, arXiv:0705.0107]

- ν_e disappearance experiments:

$$\sin^2 2\vartheta_{ee} = 4|U_{e4}|^2 \left(1 - |U_{e4}|^2\right) \simeq 4|U_{e4}|^2$$

- ν_μ disappearance experiments:

$$\sin^2 2\vartheta_{\mu\mu} = 4|U_{\mu 4}|^2 \left(1 - |U_{\mu 4}|^2\right) \simeq 4|U_{\mu 4}|^2$$

- $\nu_\mu \rightarrow \nu_e$ experiments:


$$\sin^2 2\vartheta_{e\mu} = 4|U_{e4}|^2 |U_{\mu 4}|^2 \simeq \frac{1}{4} \sin^2 2\vartheta_{ee} \sin^2 2\vartheta_{\mu\mu}$$


- Upper bounds on $\sin^2 2\vartheta_{ee}$ and $\sin^2 2\vartheta_{\mu\mu}$ \implies strong limit on $\sin^2 2\vartheta_{e\mu}$

[Okada, Yasuda, Int. J. Mod. Phys. A12 (1997) 3669-3694, arXiv:hep-ph/9606411]

[Bilenky, Giunti, Grimus, Eur. Phys. J. C1 (1998) 247, arXiv:hep-ph/9607372]

3+1 Schemes







PGoF = 0.006%

- ▶ Strong tension between $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ appearance and disappearance limits ($\bar{\nu}_e \rightarrow \bar{\nu}_e$ and mainly $\nu_\mu \rightarrow \nu_\mu$)
- ▶ Tension reduced in 3+2, 3+1+NSI [talk by T. Schwetz]
- ▶ CPT Violation?

[Barger, Marfatia, Whisnant, PLB 576 (2003) 303]

[Giunti, Laveder, PRD 82 (2010) 093016, PRD 83 (2011) 053006]

Antineutrino Oscillations in 3+1 Schemes



$$\chi^2_{\min} = 87.0$$

$$NdF = 83$$

$$GoF = 36\%$$

$$\Delta m^2 = 0.42 \text{ eV}^2 \quad \sin^2 2\vartheta_{e\mu} = 0.016 \quad \sin^2 2\vartheta_{ee} = 0.020 \quad \sin^2 2\vartheta_{\mu\mu} = 0.65$$

Prediction: large SBL $\bar{\nu}_\mu$ disappearance at $\Delta m^2 \gtrsim 0.1 \text{ eV}^2$

[Giunti, Laveder, PRD 83 (2011) 053006, arXiv:1012.0267]

Gallium Anomaly


Gallium Radioactive Source Experiments [talk by E. Bellotti]

Tests of the solar neutrino detectors **GALLEX** (Cr1, Cr2) and **SAGE** (Cr, Ar)

Detection Process: $\nu_e + {}^{71}\text{Ga} \rightarrow {}^{71}\text{Ge} + e^-$


ν_e Sources: $e^- + {}^{51}\text{Cr} \rightarrow {}^{51}\text{V} + \nu_e$ $e^- + {}^{37}\text{Ar} \rightarrow {}^{37}\text{Cl} + \nu_e$

	${}^{51}\text{Cr}$				${}^{37}\text{Ar}$	
E [keV]	747	752	427	432	811	813
B.R.	0.8163	0.0849	0.0895	0.0093	0.902	0.098



[SAGE, PRC 73 (2006) 045805, nucl-ex/0512041]


[SAGE, PRC 59 (1999) 2246, hep-ph/9803418]



[SAGE, PRC 59 (1999) 2246, hep-ph/9803418]

$$\langle L \rangle_{\text{GALLEX}} = 1.9 \text{ m}$$

$$\langle L \rangle_{\text{SAGE}} = 0.6 \text{ m}$$



[SAGE, PRC 73 (2006) 045805, nucl-ex/0512041]


$$\begin{aligned}
 R_B^{\text{Gallex-Cr1}} &= 0.953 \pm 0.11 \\
 R_B^{\text{Gallex-Cr2}} &= 0.812^{+0.10}_{-0.11} \\
 R_B^{\text{SAGE-Cr}} &= 0.95 \pm 0.12 \\
 R_B^{\text{SAGE-Ar}} &= 0.791^{+0.084}_{-0.078}
 \end{aligned}$$

$$R_B^{\text{Ga}} = 0.86 \pm 0.05$$

Bahcall cross section

Only exp uncertainties!

- Deficit could be due to overestimate of
 $\sigma(\nu_e + {}^{71}\text{Ga} \rightarrow {}^{71}\text{Ge} + e^-)$
- Calculation: Bahcall, PRC 56 (1997) 3391, hep-ph/9710491



- $\sigma_{\text{G.S.}}$ related to measured $\sigma(e^- + {}^{71}\text{Ge} \rightarrow {}^{71}\text{Ga} + \nu_e)$:

$$\sigma_{\text{G.S.}}({}^{51}\text{Cr}) = 55.3 \times 10^{-46} \text{ cm}^2 (1 \pm 0.004)_{3\sigma}$$

$$\sigma({}^{51}\text{Cr}) = \sigma_{\text{G.S.}}({}^{51}\text{Cr}) \left(1 + 0.669 \frac{\text{BGT}_{175 \text{ keV}}}{\text{BGT}_{\text{G.S.}}} + 0.220 \frac{\text{BGT}_{500 \text{ keV}}}{\text{BGT}_{\text{G.S.}}} \right)$$

- Contribution of Excited States only 5%!

► Bahcall:

[Bahcall, PRC 56 (1997) 3391, hep-ph/9710491]

from $p + {}^{71}\text{Ga} \rightarrow {}^{71}\text{Ge} + n$ measurements [Krofcheck et al., PRL 55 (1985) 1051]

$$\frac{\text{BGT}_{175\text{ keV}}}{\text{BGT}_{\text{G.S.}}} < 0.056 \Rightarrow \frac{\text{BGT}_{175\text{ keV}}}{\text{BGT}_{\text{G.S.}}} = \frac{0.056}{2} \quad \frac{\text{BGT}_{500\text{ keV}}}{\text{BGT}_{\text{G.S.}}} = 0.146$$

3σ lower limit: $\frac{\text{BGT}_{175\text{ keV}}}{\text{BGT}_{\text{G.S.}}} = \frac{\text{BGT}_{500\text{ keV}}}{\text{BGT}_{\text{G.S.}}} = 0$

3σ upper limit: $\frac{\text{BGT}_{175\text{ keV}}}{\text{BGT}_{\text{G.S.}}} < 0.056 \times 2 \quad \frac{\text{BGT}_{500\text{ keV}}}{\text{BGT}_{\text{G.S.}}} = 0.146 \times 2$

$$\sigma({}^{51}\text{Cr}) = 58.1 \times 10^{-46} \text{ cm}^2 \left(1^{+0.036}_{-0.028}\right)_{1\sigma} \Rightarrow R_B^{\text{Ga}} = 0.86 \pm 0.06$$


► Haxton:

[Hata, Haxton, PLB 353 (1995) 422, nucl-th/9503017; Haxton, PLB 431 (1998) 110, nucl-th/9804011]

"a sophisticated shell model calculation is performed ... for the transition to the first excited state in ${}^{71}\text{Ge}$. The calculation predicts destructive interference between the (p, n) spin and spin-tensor matrix elements."

$$\sigma({}^{51}\text{Cr}) = 63.9 \times 10^{-46} \text{ cm}^2 (1 \pm 0.106)_{1\sigma} \Rightarrow R_H^{\text{Ga}} = 0.78 \pm 0.13$$

- $R_H^{Ga} = 0.78 \pm 0.13$ exp and the uncertainties added in quadrature
- $R^{Ga} = R_{exp}^{Ga} / R_{the}^{Ga}$ probability distribution of ratio is not Gaussian



$$p_{R^{Ga}}(r) = \int_{R_{gs}^{Ga}}^{\infty} p_{R_{exp}^{Ga}}(rs) p_{R_{the}^{Ga}}(s) s ds$$

$$R^{Ga} = 0.76^{+0.09}_{-0.08}$$

$R^{Gallex-Cr1}$	$= 0.84^{+0.13}_{-0.12}$
$R^{Gallex-Cr2}$	$= 0.71^{+0.12}_{-0.11}$
$R^{SAGE-Cr}$	$= 0.84^{+0.14}_{-0.13}$
$R^{SAGE-Ar}$	$= 0.70^{+0.10}_{-0.09}$

[Giunti, Laveder, arXiv:1006.3244]

$$P_{\nu_e \rightarrow \nu_e}^{\text{SBL}}(L, E) = 1 - \sin^2 2\vartheta \sin^2 \left(\frac{\Delta m^2 L}{4E} \right)$$

$$R^k(\sin^2 2\vartheta, \Delta m^2) = \frac{\int_k dV L^{-2} \sum_i b_i^k \sigma_i^k P_{\nu_e \rightarrow \nu_e}^{\text{SBL}}(L, E_i)}{\sum_i b_i^k \sigma_i^k \int_k dV L^{-2}}$$


$k = \text{GALLEX-Cr1, GALLEX-Cr2, SAGE-Cr, SAGE-Ar}$

$$R^k = R_{\text{exp}}^k / R_{\text{the}}^k \quad \text{fully correlated theoretical uncertainty!}$$

$$p_{\vec{R}}(\vec{r}) = \int_{R_{\text{gs}}^k}^{\infty} \left[\prod_k p_{R_{\text{exp}}^k}(r^k s) \right] p_{R_{\text{the}}^k}(s) s^4 ds$$

$$\mathcal{L}(\sin^2 2\vartheta, \Delta m^2) = p_{\vec{R}}(\vec{R}(\sin^2 2\vartheta, \Delta m^2))$$

$$\chi^2(\sin^2 2\vartheta, \Delta m^2) = -2 \ln \mathcal{L}(\sin^2 2\vartheta, \Delta m^2) + \text{constant}$$



$$\Delta\chi^2_{\text{No Osc.}} = 9.7$$

No Osc. disfavored
at 99.23 % C.L
(2.7 σ)



Osc.

$$\sin^2 2\vartheta_{\text{bf}} = 0.51$$

$$\Delta m_{\text{bf}}^2 = 2.24 \text{ eV}^2$$

[Giunti, Laveder, arXiv:1006.3244]


Reactor Antineutrino Anomaly



- $R_{\text{rates}} = 0.946 \pm 0.024$
- Improved hint of oscillations given by Bugey energy spectrum with old reactor fluxes
- $\sin^2 2\vartheta_{\text{bf}} = 0.059$ $\Delta m_{\text{bf}}^2 = 1.89 \text{ eV}^2$

[Acero, Giunti, Laveder, PRD 78 (2008) 073009, arXiv:0711.4222]


Reactor Antineutrino Anomaly



$$\sin^2 2\vartheta_{bf} = 0.059$$

$$\Delta m_{bf}^2 = 1.89 \text{ eV}^2$$


Gallium Neutrino Anomaly



$$\sin^2 2\vartheta_{bf} = 0.51$$

$$\Delta m_{bf}^2 = 2.24 \text{ eV}^2$$

Gallium Anomaly + Reactor Anomaly



$$\chi^2_{\min} = 59.6$$

$$\text{NdF} = 71$$


$$\text{GoF} = 83\%$$

$$\sin^2 2\theta = 0.11$$

$$\Delta m^2 = 2.95 \text{ eV}^2$$


$$\text{PGoF} = 4.6\%$$

Old Reactor $\bar{\nu}_e$ Fluxes



PGoF = 2.3%


New Reactor $\bar{\nu}_e$ Fluxes




PGoF = 4.6%

Implications of Gallium and Reactor Anomalies

β Decay



$(\beta\beta)_{0\nu}$ Decay




$$m_\beta^2 = \sum_k |U_{ek}|^2 m_k^2$$

$$m_{\beta\beta} = \left| \sum_k U_{ek}^2 m_k \right|$$

[Giunti, Laveder, In Preparation]

MiniBooNE Low-Energy Anomaly




[PRL 102 (2009) 101802, arXiv:0812.2243]

Our Hypothesis: $N_{\nu,j}^{\text{the}} = f_\nu \left(P_{\nu_e \rightarrow \nu_e} N_{\nu_e,j}^{\text{cal}} + N_{\nu_\mu,j}^{\text{cal}} \right)$

[Giunti, Laveder, PRD 77 (2008) 093002, arXiv:0707.4593; PRD 80 (2009) 013005, arXiv:0902.1992]

$$N_{\nu,j}^{\text{the}} = f_\nu \left(P_{\nu_e \rightarrow \nu_e} N_{\nu_e,j}^{\text{cal}} + N_{\nu_\mu,j}^{\text{cal}} \right)$$

- ▶ Estimated 15% uncertainty of the calculated neutrino flux [MiniBooNE, PRD 79 (2009) 072002, arXiv:0806.1449] is consistent with measured ratio 1.21 ± 0.24 of detected and predicted charged-current quasi-elastic ν_μ events [MiniBooNE, PRL 100 (2008) 032301, arXiv:0706.0926]
- ▶ We fit MiniBooNE ν_e and ν_μ data using the info at http://www-boone.fnal.gov/for_physicists/data_release/lowe/



No Osc. & $f_\nu = 1$

$$\chi^2_{\min} = 14.3 + 5.4$$

$$NdF = 3 + 16$$

$$GoF = 41\%$$

Our Hypothesis

$$\chi^2_{\min} = 2.0 + 7.6$$

$$NdF = 16$$

$$GoF = 89\%$$


$$f_\nu = 1.26$$

$$\sin^2 2\vartheta = 0.32$$

$$\Delta m^2 = 1.84 \text{ eV}^2$$

[Giunti, Laveder, PRD 82 (2010) 053005, arXiv:1005.4599]

MiniBooNE + Gallium



$$\chi^2_{\min} = 2.3 + 9.2$$

$$NdF = 20$$

$$GoF = 93\%$$


$$\sin^2 2\vartheta = 0.27$$

$$\Delta m^2 = 1.92 \text{ eV}^2$$

$$PGoF = 93\%$$

[Giunti, Laveder, PRD 82 (2010) 053005, arXiv:1005.4599]



MiniBooNE + Gallium + Reactors



[Mention et al., arXiv:1101.2755]

Future

- ▶ New Gallium source experiments: ν_e disappearance [Gavrin et al, arXiv:1006.2103]
- ▶ CPT test: ν_e and $\bar{\nu}_e$ disappearance
- ▶ Beta-Beam experiments: [Antusch, Fernandez-Martinez, PLB 665 (2008) 190, arXiv:0804.2820]



- ▶ Neutrino Factory experiments: [Giunti, Laveder, Winter, PRD 80 (2009) 073005, arXiv:0907.5487]



- ▶ New ν_e and $\bar{\nu}_e$ radioactive source experiments with low-threshold neutrino elastic scattering detectors.
- ▶ Borexino: talk by M. Pallavicini

[Ianni, Montanino, Scioscia, EPJC 8 (1999) 609, arXiv:hep-ex/9901012]

- ▶ LENS (Low Energy Neutrino Spectroscopy): [Agarwalla, Raghavan, arXiv:1011.4509]



- ▶ Spherical Gaseous TPC: [Vergados, Giomataris, Novikov, arXiv:1103.5307]
 - ▶ Targets: ${}^{131}\text{Xe}$, ${}^{40}\text{Ar}$, ${}^{20}\text{Ne}$, ${}^4\text{He}$.
 - ▶ Sources: ${}^{37}\text{Ar}$, ${}^{51}\text{Cr}$, ${}^{65}\text{Zn}$, ${}^{32}\text{P}$.

Conclusions 1


- ▶ Suggestive LSND and MiniBooNE agreement on SBL $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$
- ▶ Hint in favor of sterile neutrinos is compatible with cosmological data, but mass is limited
- ▶ Two experimental tensions:
 - ▶ LSND and MiniBooNE $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ vs MiniBooNE $\nu_\mu \rightarrow \nu_e$ (CP violation?)
 - ▶ LSND and MiniBooNE $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ vs $\bar{\nu}_e$ and ν_μ disappearance limits
- ▶ CPT-invariant 3+1 Four-Neutrino Mixing is strongly disfavored (no CP violation and tension between appearance and disappearance)
- ▶ 3+2 can explain CP violation and reduce tension between appearance and disappearance with New Reactor $\bar{\nu}_e$ Fluxes [talk by T. Schwetz]
- ▶ 3+1+NSI has CP violation and reduced appearance-disappearance tension
- ▶ CPT-violating 3+1 Mixing \implies testable large SBL $\bar{\nu}_\mu$ disappearance

Conclusions 2


- ▶ Interesting possible agreement of
 - ▶ Gallium Anomaly (SBL ν_e disappearance)
 - ▶ Reactor Anomaly (SBL $\bar{\nu}_e$ disappearance)
- ▶ Testable Predictions:
 - ▶ $m_\beta \sim 0.12 - 0.71 \text{ eV}$ (2σ)
 - ▶ $m_{\beta\beta} \sim 0.011 - 0.15 \text{ eV}$ (2σ)
- ▶ Exciting experimental results, but interpretation is not clear:
 - ▶ Explanation of all data needs at least two new physical effects.
 - ▶ Without CPT violation tensions do not disappear completely.
 - ▶ Possible that some experiments are giving misleading information.
- ▶ New short-baseline neutrino oscillation experiments are needed!

Backup Slides


Four-Neutrino Schemes: 2+2 and 3+1



2+2 Four-Neutrino Schemes




"normal"



"inverted"

2+2 Schemes are strongly disfavored by solar and atmospheric data




[Maltoni, Schwetz, Tortola, Valle, New J. Phys. 6 (2004) 122, arXiv:hep-ph/0405172]

$$\eta_s = |U_{s1}|^2 + |U_{s2}|^2$$

$$99\% \text{ CL: } \begin{cases} \eta_s < 0.25 & (\text{solar} + \text{KamLAND}) \\ \eta_s > 0.75 & (\text{atmospheric} + \text{K2K}) \end{cases}$$

3+1 Four-Neutrino Schemes



Perturbation of 3-ν Mixing

$$|U_{e4}|^2 \ll 1 \quad |U_{\mu 4}|^2 \ll 1 \quad |U_{\tau 4}|^2 \ll 1 \quad |U_{s4}|^2 \simeq 1$$