


THE GROWING TOOLBOX OF PERTURBATIVE QCD

Lorenzo Magnea

University of Torino - INFN Torino

ECT* - Trento - 11/09/2015



Outline

- Introduction
- QCD at future colliders
- Selected examples of new tools
- Soft gluons beyond leading power
- Outlook

INTRODUCTION



Where we stand

Where we stand

 The **LHC** had a **wonderful first run**, culminating in the Nobel-Prize-winning **discovery of the Higgs boson**.

***** *We expected it, but it was a great achievement nonetheless.*

Where we stand

 The **LHC** had a **wonderful first run**, culminating in the Nobel-Prize-winning **discovery of the Higgs boson**.

** We expected it, but it was a great achievement nonetheless.*

 The **Standard Model** of particle physics has been **verified** to an unexpected, even **amazing degree of accuracy**.

** But the Holy Grail - new physics - is not in sight.*

Where we stand

- 📌 The **LHC** had a **wonderful first run**, culminating in the Nobel-Prize-winning **discovery of the Higgs boson**.
 - * *We expected it, but it was a great achievement nonetheless.*
- 📌 The **Standard Model** of particle physics has been **verified** to an unexpected, even **amazing degree of accuracy**.
 - * *But the Holy Grail - new physics - is not in sight.*
- 📌 The **absence** of **new physics** signals is quietly **shifting the focus** of the theory community **away from the 'Standard BSM'** scenario of 'Naturalness + Supersymmetry'.
 - * *LHC Run Two may effectively kill many cherished models.*

Where we stand

- 📌 The **LHC** had a **wonderful first run**, culminating in the Nobel-Prize-winning **discovery of the Higgs boson**.
 - * *We expected it, but it was a great achievement nonetheless.*
- 📌 The **Standard Model** of particle physics has been **verified** to an unexpected, even **amazing degree of accuracy**.
 - * *But the Holy Grail - new physics - is not in sight.*
- 📌 The **absence** of **new physics** signals is quietly **shifting the focus** of the theory community **away from the 'Standard BSM'** scenario of 'Naturalness + Supersymmetry'.
 - * *LHC Run Two may effectively kill many cherished models.*
- 📌 The **probability** of striking, **macroscopic new physics signatures** with a **moderate** increase in energy **appears low**.
 - * *We will probably have to disentangle small distortions from large SM backgrounds.*

Where we stand

- 📌 The **LHC** had a **wonderful first run**, culminating in the Nobel-Prize-winning **discovery of the Higgs boson**.
 - * *We expected it, but it was a great achievement nonetheless.*
- 📌 The **Standard Model** of particle physics has been **verified** to an unexpected, even **amazing degree of accuracy**.
 - * *But the Holy Grail - new physics - is not in sight.*
- 📌 The **absence** of **new physics** signals is quietly **shifting the focus** of the theory community **away from the 'Standard BSM'** scenario of 'Naturalness + Supersymmetry'.
 - * *LHC Run Two may effectively kill many cherished models.*
- 📌 The **probability** of striking, **macroscopic new physics signatures** with a **moderate** increase in energy **appears low**.
 - * *We will probably have to disentangle small distortions from large SM backgrounds.*
- 📌 In the **coming years** the need will be for **precise predictions** of **increasingly complex** SM processes.
 - * *A great time for the QCD community.*

Where we stand

- 🎤 The **LHC** had a **wonderful first run**, culminating in the Nobel-Prize-winning **discovery of the Higgs boson**.
 - * *We expected it, but it was a great achievement nonetheless.*
- 🎤 The **Standard Model** of particle physics has been **verified** to an unexpected, even **amazing degree of accuracy**.
 - * *But the Holy Grail - new physics - is not in sight.*
- 🎤 The **absence** of **new physics** signals is quietly **shifting the focus** of the theory community **away from the 'Standard BSM'** scenario of 'Naturalness + Supersymmetry'.
 - * *LHC Run Two may effectively kill many cherished models.*
- 🎤 The **probability** of striking, **macroscopic new physics signatures** with a **moderate** increase in energy **appears low**.
 - * *We will probably have to disentangle small distortions from large SM backgrounds.*
- 🎤 In the **coming years** the need will be for **precise predictions** of **increasingly complex** SM processes.
 - * *A great time for the QCD community.*
- 🎤 The **run-up** to the LHC has seen a **vast effort** and **great progress** in **precision phenomenology**: PDF's, jets, hard cross sections, resummations, and more.
 - * *A continuing effort that will hopefully pay off during Run Two!*

QCD BEYOND LHC



QCD at future colliders

QCD at future colliders

Hadron colliders:

QCD at future colliders

Hadron colliders:



Hadron Collider
you build,
much QCD
you need.

QCD at future colliders

Lepton colliders:


QCD at future colliders

Lepton colliders:




Lepton Collider
you build,
much QCD
you need.

Lepton collider jets




ttH productions can yield 8-jet final states



“Underlying event” at CLIC3TeV (Vos, at LCWS14)

- Relatively **small samples** at lepton colliders
 - ◆ *Important to use hadronic final states.*
- Interesting processes yield **many jets**.
 - ◆ *The best available jet tool are needed.*
- There **actually is** an **Underlying Event**.
 - ◆ *$\gamma\gamma \rightarrow$ hadrons, pair production.*
- Jet algorithms** are important.
 - ◆ *Boost invariance is less relevant.*
- Algorithms affect jet **shapes** and **areas**.
 - ◆ *Forward region has high backgrounds.*
- Jet **radius** analysis must be **re-tuned**.
 - ◆ *Strong energy dependence of UE.*
- Jet **substructure** analysis still important.
 - ◆ *But fatter jets and unboosted objects.*

Standard model parameters




Threshold $t\bar{t}$ production at N^3 LO (Beneke et al.)

- M_{top} must be **precisely defined**.
 - ◆ *Minimize non-perturbative effects.*
- **Lepton colliders** provide the best precision.
 - ◆ *Scanning the threshold for $t\bar{t}$ production.*
- The **most refined** tools of **QCD** are needed.
 - ◆ *High orders, effective theories, resummations.*
- M_{top} can be **determined** to within **50 MeV**.

- Unsolved **discrepancies** in event shape **α_s fits**.
 - ◆ *Unexpected influence of power corrections.*

$$\begin{aligned} \alpha_s(M_Z^2) &= 0.1172 \pm 0.0022 && \text{thrust (BS)} \\ \alpha_s(M_Z^2) &= 0.1220 \pm 0.0031 && \text{jet mass (SC)} \\ \alpha_s(M_Z^2) &= 0.1135 \pm 0.0010 && \text{thrust (AFHMS)} \end{aligned}$$

- **Very accurate** studies: **NNLO, N^3 LL, PC's**.
 - ◆ *Do we need a more precise definition of α_s ?*




Do we really know α_s as well as we think? (Mateu 2013)

NEW TOOLS



Two loops is the new one loop


- **Two-loop** calculations are **not yet** a **commodity**: they are largely **custom-made** and **expensive**.
- A major **stumbling block** has been the **subtraction** of **infrared** and **collinear** singularities.
- **Progress** has been slow but is rapidly **speeding up**: **automation** is on the way.




From **Claude Duhr's** talk at **ICHEP 2014**

Two loops is the new one loop

- Two-loop calculations are **not yet** a **commodity**: they are largely **custom-made** and **expensive**.
- A major **stumbling block** has been the **subtraction** of **infrared** and **collinear** singularities.
- Progress** has been slow but is rapidly **speeding up**: **automation** is on the way.




V. Smirnov hep-ph/9905323




Two loops is the new one loop

- Two-loop calculations are **not yet** a **commodity**: they are largely **custom-made** and **expensive**.
- A major **stumbling block** has been the **subtraction** of **infrared** and **collinear** singularities.
- Progress** has been slow but is rapidly **speeding up**: **automation** is on the way.



From **Claude Duhr's** talk at **ICHEP 2014**

V. Smirnov [hep-ph/9905323](https://arxiv.org/abs/hep-ph/9905323)






Full color
Masses

The NNLO subtraction problem

- A **well-known** problem: **infrared** and **collinear** divergences **cancel** between final states with different **particle content** and different **phase spaces**.
- The **cancellation** must be **performed locally** in phase space to allow for **generic observables**.
- “**Simple**” subtraction **counterterms** must be constructed in **each** phase space.
- A **surprisingly hard problem**, on the table for more than a **decade**.

$$d\hat{\sigma}_{NNLO} \sim \int_{d\Phi_{m+2}} d\hat{\sigma}_{NNLO}^{RR} + \int_{d\Phi_{m+1}} d\hat{\sigma}_{NNLO}^{RV} + \int_{d\Phi_m} d\hat{\sigma}_{NNLO}^{VV}$$

Different final-state multiplicities
conspire to cancel infrared and
collinear poles



	analytic	FS colour	IS colour	local
antenna subtraction	✓	✓	✓	✗
STRIPPER	✗	✓	✓	✓
q_T subtraction	✓	✗	✓	✓
reverse unitarity	✓	✗	✓	-
Trócsányi et al	✗	✓	✗	✓

Comparing subtraction algorithms, from James Currie's talk at LoopFest

- Several** solutions are now **available**.
- Analytical** vs. **numerical** approaches.
- Dedicated** vs. **general** algorithms.
- Several groups** at work.
- New: N-jettiness** subtraction.
- No silver bullet** yet.

Iterated integrals

- A **large class** of integrals arising from Feynman diagrams (**but not all!**) can be expressed as “**iterated integrals**”, yielding functions in the **class of polylogarithms**. At **one** loop

$$\log z = - \int_0^{1-z} \frac{dt}{1-t}, \quad \text{Li}_2(z) = \int_0^z \frac{dt}{t} \int_0^t \frac{du}{1-u}$$

- At **higher orders** one encounters **more general** examples, such as **Harmonic Polylogarithms** or **Goncharov Polylogarithms**

$$G_{a_1, \dots, a_n}(z) \equiv \int_0^z \frac{dt}{t - a_1} G_{a_2, \dots, a_n}(t),$$

- Notice** that **all** these integrals are of a “**d log**” form: at each step one integrates **over the logarithm** of a simple (here linear) function of the integration variables.
- The **parameters** a_n are the **locations** of **singular** points and have **physical meaning**.
- Iterated integrals** are organized by a **powerful underlying algebraic structure**, described by the “**Symbol**” map or by a Hopf algebra with a notion of “**Co-product**” (**Duhr**).
- In particular **each** such function can be assigned a “**weight**” w , equal to the **number of iterations**. For example $\text{Li}_2(z)$ has weight $w = 2$, and $\zeta(n)$ has weight $w = n$.
- These structures were **uncovered** in the context of studies of **N=4 Super Yang-Mills theory** amplitudes, where they have played a **pivotal role**.
- We **now** see **powerful** new **applications** to ordinary **QCD** (**Henn, Smirnov, Von Manteuffel**)

A good N^PLO harvest



- A wealth of results for on-shell production of electroweak boson pairs: ZZ , W^+W^- , $W\gamma$, $Z\gamma$ (Zurich group, see e.g. 1405.2219, 1507.062570).
- All helicity amplitudes for $pp \rightarrow VV'$ at two-loop now known. (Caola, Henn, Melnikov, Smirnov², 1404.5590, 1408.6409, 1503.08759).
- Preliminary results for $\gamma^* \gamma^*$ production (Anastasiou *et al.* 1408.4546).
- Differential distributions for associated ZH production (Ferrera, Grazzini, Tramontano, 1407.4747).
- Differential distributions in Higgs + one jet production (Chen *et al.* 1408.5325; Boughezal *et al.* 1504.07922).
- Differential distributions in W,Z + one jet production (Boughezal *et al.* 1504.02131; Gehrmann *et al.* 1507.02850;).
- Differential distributions for t -channel single top production in the 'structure function' approximation (Brucherseifer, Caola, Melnikov, 1404.7116).
- Progress towards the construction of a general basis for two-loop master integrals (Mastrolia *et al.*; Badger, Frellesvig, Zhang, 1407.3133).
- The three-loop angle-dependent cusp anomalous dimension in QCD (Grozin, Henn, Korchemsky, Marquard, 1409.0023).
- The three-loop massless multi-parton soft anomalous dimension in QCD (Almelid, Duhr, Gardi 1507.00047).


Three loops is the new two loops

- After the **landmark** calculation of **three-loop DIS structure functions** by Moch, Vermaseren and Vogt a decade ago, the next **PQCD challenge** has been the computation of a **cross section without an OPE** at three loops. The “Drell-Yan” process is the best candidate.
- At **LHC**, “Drell-Yan” means **vector boson production** and **Higgs production via gluon fusion**. The phenomenological **impact** is **evident**, especially given the **large corrections** to **Higgs** production at one and two loops.
- Approximate** three-loop results using **threshold** and **Regge** limits **exist** (Moch, Vogt, 2005; LM, Laenen, 2005; Ball, Bonvini, Forte, Marzani, Ridolfi, 2013).
- The **full calculation** is now **complete** (Anastasiou, Duhr, Dulat, Furlan, Gehrmann, Herzog, Mistiberg 1503.06056), in a **threshold expansion** to essentially arbitrary order.
- The **threshold** expansion is also a “**soft gluon**” expansion

$$\hat{\sigma} = \hat{\sigma}(z), \quad z = \frac{Q^2}{\hat{s}}, \quad \hat{\sigma}(z) = \hat{\sigma}_{SV} + \hat{\sigma}_0 + (1-z)\hat{\sigma}_1 + \mathcal{O}[(1-z)^2]$$

- The **three-loop soft-virtual** contribution is **fully predicted** by threshold **resummation** **except** for $\delta(1-z)$ contributions. Resummation at **next-to-leading power** is under study.
- The “Drell-Yan” **timeline**: **1979 - 1991 - 2002 - 2015**.

Three loops is the new two loops



The partonic cross section for SM Higgs production through gluon fusion up to N³LO, with scale uncertainties.


Bold projections

Bold projections

What could be achieved on a **ten-year** time scale, if the **community** is sufficiently **motivated** to keep up this **LHC-driven** energetic work, and taking into account **hardware progress**?


Bold projections

What could be achieved on a **ten-year** time scale, if the **community** is sufficiently **motivated** to keep up this **LHC-driven** energetic work, and taking into account **hardware progress**?



Bold projections


What could be achieved on a **ten-year** time scale, if the **community** is sufficiently **motivated** to keep up this **LHC-driven** energetic work, and taking into account **hardware progress**?



- Deep Inelastic Scattering (inclusive).
 - ◆ *Structure functions at four loops, N^4LL threshold resummation, detailed analytic power corrections.*
 - ◆ *Significant impact on parton distributions.*
- Electron-positron annihilation (inclusive).
 - ◆ *Event shapes and at three loops, N^4LL threshold resummation, detailed analytic power correction.*
 - ◆ *Significant impact on α_s .*
- Hadron-hadron colliders (inclusive).
 - ◆ *Electroweak annihilation at four loops and N^4LL .*
- Hadron-hadron colliders (exclusive).
 - ◆ *Fully differential automated distributions at NNLO.*
 - ◆ *Few-jet cross section at three loops.*
 - ◆ *Control of threshold and non-threshold logarithms.*

Bold projections


What could be achieved on a **ten-year** time scale, if the **community** is sufficiently **motivated** to keep up this **LHC-driven** energetic work, and taking into account **hardware progress**?



- Deep Inelastic Scattering (inclusive).
 - ◆ *Structure functions at four loops, N^4LL threshold resummation, detailed analytic power corrections.*
 - ◆ *Significant impact on parton distributions.*
- Electron-positron annihilation (inclusive).
 - ◆ *Event shapes and at three loops, N^4LL threshold resummation, detailed analytic power correction.*
 - ◆ *Significant impact on α_s .*
- Hadron-hadron colliders (inclusive).
 - ◆ *Electroweak annihilation at four loops and N^4LL .*
- Hadron-hadron colliders (exclusive).
 - ◆ *Fully differential automated distributions at NNLO.*
 - ◆ *Few-jet cross section at three loops.*
 - ◆ *Control of threshold and non-threshold logarithms.*
- **Full control** of SM phenomenology **below percent level.**

Bold projections

What could be achieved on a **ten-year** time scale, if the **community** is sufficiently **motivated** to keep up this **LHC-driven** energetic work, and taking into account **hardware progress**?



- Deep Inelastic Scattering (inclusive).
 - ◆ *Structure functions at four loops, N^4LL threshold resummation, detailed analytic power corrections.*
 - ◆ *Significant impact on parton distributions.*
- Electron-positron annihilation (inclusive).
 - ◆ *Event shapes and at three loops, N^4LL threshold resummation, detailed analytic power correction.*
 - ◆ *Significant impact on α_s .*
- Hadron-hadron colliders (inclusive).
 - ◆ *Electroweak annihilation at four loops and N^4LL .*
- Hadron-hadron colliders (exclusive).
 - ◆ *Fully differential automated distributions at NNLO.*
 - ◆ *Few-jet cross section at three loops.*
 - ◆ *Control of threshold and non-threshold logarithms.*
- **Full control** of SM phenomenology **below percent level**.
 - ◆ *We must hope we will NOT need this.*

THRESHOLDS



Logarithms

📌 **Multi-scale** problems in **renormalizable** quantum field theories have perturbative corrections of the form $\alpha_s^n \log^k (Q_i^2/Q_j^2)$, which may **spoil** the reliability of the perturbative expansion. However, they **carry important physical information**.

- **Renormalization** and **factorization** logs: $\alpha_s^n \log^n (Q^2/\mu^2)$
- **High-energy logs**: $\alpha_s^n \log^{n-1} (s/t)$
- **Sudakov** logs: $\alpha_s^n \log^{2n-1} (1-z)$, $1-z = W^2/Q^2, 1-M^2/\hat{s}, Q_\perp^2/Q^2, \dots$

📌 **Logarithms** encode **process-independent** features of perturbation theory. For **Sudakov** logs: the structure of **infrared** and **collinear** divergences.

$$\underbrace{\frac{1}{\epsilon}}_{\text{virtual}} + \underbrace{(Q^2)^\epsilon \int_0^{m^2} \frac{dk^2}{(k^2)^{1+\epsilon}}}_{\text{real}} \implies \ln(m^2/Q^2)$$

- For **inclusive** observables: **analytic** resummation to high logarithmic accuracy.
- For **exclusive** final states: **parton shower** event generators, **(N(N))LL** accuracy. 


Gradually merging

📌 **Resummation** probes the **all-order structure** of perturbation theory.

- **Non-perturbative** contributions to QCD cross sections can be estimated.
- Links to the **strong coupling** regime can be established for special gauge theories.

Logarithms at work

Predictions for the thrust distribution at LEP (E. Gardi, J. Rathsmann, 02)




The thrust distribution is computed with NLL soft gluon resummations and with various models of power correction (shape functions). For illustrative purposes: many improvements in later work.

Logarithms at work

Predictions for the thrust distribution at LEP (E. Gardi, J. Rathsmann, 02)


Note shift in the distribution due to non-perturbative corrections extrapolated from all-order resummed result



The thrust distribution is computed with NLL soft gluon resummations and with various models of power correction (shape functions). For illustrative purposes: many improvements in later work.

Logarithms at work


Predictions for the Higgs boson q_T spectrum at LHC (M. Grazzini, 05)



Predictions for the q_T spectrum of Higgs bosons produced via gluon fusion at the LHC, with and without resummation, and theoretical uncertainty band of the resummed prediction.

Logarithms at work


Predictions for the Higgs boson q_T spectrum at LHC (M. Grazzini, 05)



Predictions for the q_T spectrum of Higgs bosons produced via gluon fusion at the LHC, with and without resummation, and theoretical uncertainty band of the resummed prediction.

Logarithms at work

Higgs p_T distribution somewhat shifted to higher transverse momentum but of course statistically limited.



Preliminary ATLAS data on the Higgs p_T distribution from Run I
(shown by Joey Huston at Radcor-Loopfest 2015)

The perturbative exponent

A classic way to **organize** Sudakov logarithms in terms of the **Mellin (Laplace) transform** of the momentum space cross section (**Catani et al. 93**) is to write

$$\begin{aligned} d\sigma(\alpha_s, N) &= \sum_{n=0}^{\infty} \left(\frac{\alpha_s}{\pi}\right)^n \sum_{k=0}^{2n} c_{nk} \log^k N + \mathcal{O}(1/N) \\ &= H(\alpha_s) \exp \left[\log N g_1(\alpha_s \log N) + g_2(\alpha_s \log N) + \alpha_s g_3(\alpha_s \log N) + \dots \right] + \mathcal{O}(1/N) \end{aligned}$$

This displays the main **features of Sudakov resummation**

- Predictive:** a **k**-loop calculation determines **g_k** and thus a whole **tower** of logarithms to all orders in perturbation theory.
- Effective:**
 - the **range of applicability** of perturbation theory is **extended** (finite order: **α_s log²N** small. NLL resummed: **α_s** small);
 - the renormalization **scale dependence** is naturally **reduced**.
- Theoretically interesting:** resummation **ambiguities** related to the **Landau pole** give access to non-perturbative **power-suppressed corrections**.
- Well understood:**
 - NLL** Sudakov resummations **exist** for most **inclusive** observables at hadron colliders, **NNLL** and approximate **N³LL** in simple cases.

BEYOND THRESHOLD

With D. Bonocore, E. Laenen, L. Vernazza and C. White



More logarithms

- **Threshold logarithms** are associated with kinematic variables ξ that **vanish** at **Born level** and get **corrections** that are **enhanced** because **phase space** for real radiation is **restricted** near **partonic** threshold: examples are $1 - T$, $1 - M^2/\hat{s}$, $1 - x_{BJ}$.
- At **leading power** in the threshold variable ξ logarithms are **directly related** to **soft** and **collinear divergences**: real radiation is proportional to factors of

$$\frac{1}{\xi^{1+p\epsilon}} = -\frac{1}{p\epsilon} \delta(\xi) + \left(\frac{1}{\xi}\right)_+ - p\epsilon \left(\frac{\log \xi}{\xi}\right)_+ + \dots$$

Cancels virtual IR poles

Leading power threshold logs

- **Beyond** the **leading power**, $1/\xi$, the perturbative cross section takes the form

$$\frac{d\sigma}{d\xi} = \sum_{n=0}^{\infty} \left(\frac{\alpha_s}{\pi}\right)^n \sum_{m=0}^{2n-1} \left[c_{nm}^{(-1)} \left(\frac{\log^m \xi}{\xi}\right)_+ + c_n^{(\delta)} \delta(\xi) + c_{nm}^{(0)} \log^m \xi + \dots \right]$$

Resummed to high accuracy


All-order structure in some cases

NLP threshold logs

- The **structure** of **NLP** threshold logarithms may be understood to **all orders**.

The LBKD Theorem

The **earliest evidence** that infrared effects can be **controlled** at **NLP** is **Low's theorem** (Low 58)




A radiative matrix element

$$M_\mu = e \left(\frac{p_{1\mu}'}{p_1' \cdot k} - \frac{p_{1\mu}}{p_1 \cdot k} \right) T(\nu, \Delta) + e \left(\frac{p_{1\mu}' p_{2\mu}' \cdot k}{p_1' \cdot k} - p_{2\mu}' + \frac{p_{1\mu} p_{2\mu} \cdot k}{p_1 \cdot k} - p_{2\mu} \right) \frac{\partial T(\nu, \Delta)}{\partial \nu} + O(k),$$

Low's original expression for the radiative matrix element

The LBKD Theorem

The **earliest evidence** that infrared effects can be **controlled** at **NLP** is **Low's theorem** (Low 58)



A radiative matrix element

$$M_\mu = e \left(\frac{p_{1\mu}'}{p_1' \cdot k} - \frac{p_{1\mu}}{p_1 \cdot k} \right) T(\nu, \Delta) + e \left(\frac{p_{1\mu}' p_{2\mu}' \cdot k}{p_1' \cdot k} - p_{2\mu}' + \frac{p_{1\mu} p_{2\mu} \cdot k}{p_1 \cdot k} - p_{2\mu} \right) \frac{\partial T(\nu, \Delta)}{\partial \nu} + O(k),$$


Eikonal approximation

Low's original expression for the radiative matrix element

Next-to-eikonal contribution

The LBKD Theorem

The **earliest evidence** that infrared effects can be **controlled** at **NLP** is **Low's theorem** (Low 58)



A radiative matrix element

$$M_\mu = e \left(\frac{p_{1\mu}'}{p_1' \cdot k} - \frac{p_{1\mu}}{p_1 \cdot k} \right) T(\nu, \Delta) + e \left(\frac{p_{1\mu}' p_{2\mu}' \cdot k}{p_1' \cdot k} - p_{2\mu}' + \frac{p_{1\mu} p_{2\mu} \cdot k}{p_1 \cdot k} - p_{2\mu} \right) \frac{\partial T(\nu, \Delta)}{\partial \nu} + O(k),$$

Eikonal approximation

Low's original expression for the radiative matrix element

Next-to-eikonal contribution

The **radiative matrix element** for the emission of a (next-to-) soft photon is **determined** by the **Born amplitude** T and its **first derivative** with respect to external momenta.

- Low's result established for a **single charged scalar** particle, follows from **gauge invariance**.
- It **generalizes** the well known properties of soft emissions in the **eikonal approximation**.
- The theorem was **extended** by (Burnett, Kroll 68) to particles with **spin**.
- The **LBK** theorem applies to **massive particles** and uses the **mass** as a **collinear cutoff**.
- It was **extended** to **massless** particles by (Del Duca 90), as discussed **below**.

Towards systematics

The problem of **NLP** threshold logarithms has been of interest for a **long time**, and several **different approaches** have been proposed. Recent years have seen a **resurgence of interest**, both from a **theoretical** point of view and for **phenomenology**.

- 🎤 **Early attempts** include a study of the impact of **NLP** logs on the **Higgs** cross section by **Kraemer, Laenen, Spira (98)**; work on **F_L** by **Akhoury and Sterman (99)** (logs without plus distributions are however **leading**) and work by **Grunberg et al. (07-09)** on **DIS**.
- 🎤 **Important results** can be obtained by using **physical kernels** (**Vogt et al. 09-14**) which are conjectured to be **single-logarithmic** at large **z** , which poses **constraints** on their **factorized** expression. Note in particular a **recent application** to **Higgs** production by **De Florian, Mazzitelli, Moch, Vogt (14)**.
- 🎤 **Useful approximations** can be obtained by combining **constraints** from **large N** with **high-energy** constraints for **$N \sim 1$** and **analyticity** (**Ball, Bonvini, Forte, Marzani, Ridolfi, 13**), together with **phase space** refinements.
- 🎤 **SCET techniques** can be applied and indeed may be **well-suited** to the problem: a thorough **one-loop** analysis was given in (**Larkoski, Neill, Stewart, 15**).
- 🎤 A lot of recent **formal work** on the behavior of **gauge** and **gravity** scattering **amplitudes beyond the eikonal** limit was triggered by a link to **asymptotic symmetries** of the **S** matrix (**many authors from A(ndy Strominger) to Z(vi Bern), 14-15**).

Beyond the eikonal

The **soft expansion** can be **organized beyond leading power** using either path integral techniques (Laenen, Stavenga, White 08) or diagrammatic techniques (Laenen, LM, Stavenga, White 10). The basic **idea** is **simple**, but the combinatorics **cumbersome**. For **spinors**

$$\frac{\not{p} + \not{k}}{2p \cdot k + k^2} \gamma^\mu u(p) = \left[\frac{p^\mu}{p \cdot k} + \frac{\not{k} \gamma^\mu}{2p \cdot k} - k^2 \frac{p^\mu}{2(p \cdot k)^2} \right] u(p) + \mathcal{O}(k)$$



Eikonal



NE, spin-dependent




NE, spin-independent

- A class of **factorizable** contributions **exponentiate** via **NE webs**

$$\mathcal{M} = \mathcal{M}_0 \exp \left[\sum_{D_{\text{eik}}} \tilde{C}(D_{\text{eik}}) \mathcal{F}(D_{\text{eik}}) + \sum_{D_{\text{NE}}} \tilde{C}(D_{\text{NE}}) \mathcal{F}(D_{\text{NE}}) \right].$$

- **Feynman rules** exist for the **NE** exponent, including “**seagull**” vertices.

$$\mathcal{M} = \mathcal{M}_0 \exp [\mathcal{M}_{\text{eik}} + \mathcal{M}_{\text{NE}}] (1 + \mathcal{M}_r) + \mathcal{O}(\text{NNE}).$$




A next-to-eikonal web

- **Non-factorizable** contributions involve **single gluon emission** from inside the **hard function**, and must be studied using **LBDK's theorem**.

A collinear problem

Non-factorizable contributions start at **NNLO**. For **massive** particles they can be traced to the **original LBK** theorem. For **massless** particles a **new contribution** to **NLP** logs **emerges**.




A Feynman diagram containing a collinear enhancement

- Gluon k_2 is **always** (next-to) **soft** for **EW** annihilation **near threshold**.
- When k_1 is (next-to) **soft** all logs are **captured** by **NE** rules.
- Contributions with k_1 **hard** and **collinear** are **missed** by the soft expansion.
- The **collinear pole** interferes with **soft emission** and generates **NLP** logs.
- The problem **first arises** at **NNLO**

- These contributions are **missed** by the **LBK** theorem: it applies to an **expansion** in E_k/m .
- They can be **analyzed** using the **method of regions**: the relevant **factor** is $(p \cdot k_2)^{-\epsilon}/\epsilon$.
- They **cause** the **breakdown** of **next-to-soft theorems** for amplitudes **beyond tree level**.
 ➔ the **soft** expansion and the limit $\epsilon \rightarrow 0$ **do not commute**.
- They **require** an **extension** of **LBK** to $m^2/Q < E_k < m$. It was **provided** by **Del Duca (90)**.

NLP factorization: a new jet



Soft radiation can arise either from the jets or from the hard function

$$\mathcal{A}_\mu \epsilon^\mu(k) = \mathcal{A}_\mu^J \epsilon^\mu(k) + \mathcal{A}_\mu^H \epsilon^\mu(k),$$

The amplitude for emission from the jets can be precisely defined in terms of a new jet function

$$\mathcal{A}_\mu^J = \sum_{i=1}^2 H(p_i - k; p_j, n_j) J_\mu(p_i, k, n_i) \prod_{j \neq i} J(p_j, n_j) \equiv \sum_{i=1}^2 \mathcal{A}_\mu^{J_i}.$$

Factorized contributions to the radiative amplitude

$$J_\mu(p, n, k, \alpha_s(\mu^2), \epsilon) u(p) = \int d^d y e^{-i(p-k) \cdot y} \langle 0 | \Phi_n(y, \infty) \psi(y) j_\mu(0) | p \rangle,$$

defines the radiative jet.

- At tree level the radiative jet displays the expected dependence on spin.
- Dependence on the gauge vector n^μ starts at loop level: simplifications arise for $n^2 = 0$.

$$\begin{aligned} J^{\nu(0)}(p, n, k) &= \frac{\not{k} \gamma^\nu}{2p \cdot k} - \frac{p^\nu}{p \cdot k} \\ &= -\frac{p^\nu}{p \cdot k} + \frac{k^\nu}{2p \cdot k} - \frac{i k_\alpha \Sigma^{\alpha\nu}}{2p \cdot k}. \end{aligned}$$

Beyond Low's theorem

A **slightly modified** version of **Del Duca's** result gives the **radiative amplitude** in terms of the **non-radiative** one, its **derivatives**, and the **two "jet"** functions.

$$\mathcal{A}^\mu(p_j, k) = \sum_{i=1}^2 \left\{ q_i \left(\frac{(2p_i - k)^\mu}{2p_i \cdot k - k^2} + G_i^{\nu\mu} \frac{\partial}{\partial p_i^\nu} \right) + G_i^{\nu\mu} \left[\frac{J_\nu(p_i, k, n_i)}{J(p_i, n_i)} - q_i \frac{\partial}{\partial p_i^\nu} \left(\ln J(p_i, n_i) \right) \right] \right\} \mathcal{A}(p_i; p_j).$$

The tensors $G^{\mu\nu}$ **project out the eikonal** contribution present in the first term.

$$\eta^{\mu\nu} = G^{\mu\nu} + K^{\mu\nu}, \quad K^{\mu\nu}(p; k) = \frac{(2p - k)^\nu}{2p \cdot k - k^2} k^\mu,$$

The **factorized** expression for the **radiative** amplitude can be **simplified**.


- The **jet factor** is **RG invariant**: it can be computed in **bare** perturbation theory.
- With this choice one can use that $J(p, n) = 1$ for $n^2 = 0$: it is a **pure counterterm**.
- The **choice of reference vectors** is then **physically motivated** (and **confirmed** by a complete analysis using the **method of regions**): we take $n_1 = p_2$ and $n_2 = p_1$.

$$\mathcal{A}^\mu(p_j, k) = \sum_{i=1}^2 \left(q_i \frac{(2p_i - k)^\mu}{2p_i \cdot k - k^2} + q_i G_i^{\nu\mu} \frac{\partial}{\partial p_i^\nu} + G_i^{\nu\mu} J_\nu(p_i, k) \right) \mathcal{A}(p_i; p_j).$$

For **general amplitudes**, a **full subtraction** of the residual n dependence should be **aimed at**.

Real-virtual two-loop Drell-Yan

Real-virtual corrections to EW annihilation processes involve non-factorizable contributions. NE rules cannot reproduce the perturbative result at NLP, due to collinear interference.



Real-virtual Feynman diagrams for the abelian part of the NNLO K-factor.

As a test of the LBDK factorization, we computed the C_F^2 part of the real-virtual K-factor at NNLO from ordinary Feynman diagrams, and then using the radiative amplitude integrated over phase space. As expected, plus distributions arise from the eikonal approximation, fully determined by the dressed non-radiative amplitude. Derivative terms and the projected radiative jet contribute at NLP.

- All NLP terms are correctly reproduced, including those with no logarithms.
- The radiative jet reproduces exactly the NLP collinear contribution derived by the method of regions.

$$K_{\text{rv}}^{(2)}(z) = \left(\frac{\alpha_s}{4\pi} C_F\right)^2 \left\{ \frac{32}{\epsilon^3} [\mathcal{D}_0(z) - 1] + \frac{16}{\epsilon^2} [-4\mathcal{D}_1(z) + 3\mathcal{D}_0(z) + 4L(z) - 6] \right. \\ \left. + \frac{4}{\epsilon} [16\mathcal{D}_2(z) - 24\mathcal{D}_1(z) + 32\mathcal{D}_0(z) - 16L^2(z) + 52L(z) - 49] \right. \\ \left. - \frac{128}{3}\mathcal{D}_3(z) + 96\mathcal{D}_2(z) - 256\mathcal{D}_1(z) + 256\mathcal{D}_0(z) \right. \\ \left. + \frac{128}{3}L^3(z) - 232L^2(z) + 412L(z) - 408 \right\},$$

The abelian part of the NNLO K-factor from real-virtual diagrams, omitting constants

OUTLOOK



A Perspective

- **Perturbative QCD** will play a central role in any future high-energy collider.
- Through the **vast** and **remarkable** effort of an **entire community**, ubiquitous and consistent **NLO phenomenology** at LHC is a **reality**.
- An **ongoing NNLO (r)evolution** makes it possible to imagine that in a few years we will be able to **state the same** about **NNLO phenomenology**.
- **N³LO** is the **new NNLO**.
- At some point we will have to stop ... and just resum ...
- **Leading power** threshold resummation is **highly developed** and provides some of the **most precise** predictions in perturbative **QCD**.
- **Low's theorem** is the first of **many hints** that **NLP logs** can be understood and **organized**.
- **Hard collinear** emissions **spoil** Low's theorem: a **new** radiative **jet function** emerges.
- A **complete treatment** of **NLP threshold logs** is **at hand**.
- **Much work to do** to organize a true **resummation** formula, even for **EW** annihilation: we have a more **intricate "factorization"**, we must make sure to control **double countings**.
- In order to achieve **complete generality**, we will need to include **final state jets**.

THANK YOU!